

# EXISTENCE AND MULTIPLICITY RESULTS FOR PUCCI'S OPERATORS INVOLVING NONLINEARITIES WITH ZEROS

SALOMÓN ALARCÓN<sup>†</sup>, LEONELO ITURRIAGA<sup>†,‡</sup>, AND ALEXANDER QUAAS<sup>†</sup>

<sup>†</sup> Departamento de Matemática, Universidad Técnica Federico Santa María Avenida España 1680, Casilla 110-V, Valparaíso, Chile

<sup>‡</sup> Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica, Chile.

ABSTRACT. We study the problem

$$\begin{cases} -\varepsilon^2 \mathcal{M}_{\lambda, \Lambda}^+(D^2 u) = f(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  is a smooth bounded domain in  $\mathbb{R}^N$ ,  $N > 2$ , and show it possesses nontrivial solutions for small values of  $\varepsilon$  provided  $f$  is a nonnegative continuous function which has a positive zero. The multiplicity result is based on degree theory together with a new Liouville type theorem for  $-\mathcal{M}_{\lambda, \Lambda}^+(D^2 u) = f(u)$  in  $\mathbb{R}^N$  for nonnegative nonlinearities with zeros.

## 1. INTRODUCTION

In last years an increasing interest in the study of non-proper fully nonlinear elliptic equations has arisen. This has been motivated by the current classical viscosity solution framework that started in [11], see also [12], and combined with a very rich knowledge of more simple equations that, for example, involving the Laplacian. In particular, existence of viscosity solutions to fully nonlinear elliptic equations has been extensively investigated in last twenty years for proper equations, most of them through adaptations of Perron's method, see for instance [8, 12].

In the context of non-proper fully nonlinear equations, most results have been obtained through Leray-Schauder degree theory, see [14, 5, 30, 10, 9, 17, 1]. One of the crucial points to use degree theory is to establish a priori bounds. The most classical method for obtaining a priori bounds is to use the blow-up method introduced by Gidas and Spruck, see [21], and then a Liouville type theorem in all space or in a half space. In this work, we use this kind of arguments together with a truncation procedure in order to obtain an unstable solution whereas a stable solution is obtained by the sub- and supersolutions method. Moreover, here we prove that the stable solution is isolated, from where we deduce a multiplicity result.

Before continuing, we mention that to the best of our knowledge, this paper is a first effort towards the study of singular perturbed equations in a fully nonlinear setting, and in the obtaining some asymptotic behavior of the solution. We recall that early works on singular perturbed problems are due to Ni and Takagi [27, 28], and since then this topic has become one of the most active research fields in partial differential equations. We also emphasize that most of the results in singular perturbations use strongly the variational structure of the equation, which is not present in our context.

Specifically, this paper deals with positive viscosity solutions to the problem

$$(P_\varepsilon) \quad \begin{cases} -\varepsilon^2 \mathcal{M}_{\lambda, \Lambda}^+(D^2 u) = f(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  is a bounded domain in  $\mathbb{R}^N$  with smooth boundary,  $N > 2$ ,  $\varepsilon \neq 0$  is a parameter and  $f$  satisfies  $f(x, 0) = f(x, 1) = 0$ ,  $f(x, t) > 0$  for any  $t \notin \{0, 1\}$ . Here, for fixed  $0 < \lambda \leq \Lambda$ , the

Pucci's extremal operators are defined, as in [6], by:

$$\mathcal{M}_{\lambda,\Lambda}^-(A) = \lambda \sum_{e_i > 0} e_i + \Lambda \sum_{e_i < 0} e_i \quad \text{and} \quad \mathcal{M}_{\lambda,\Lambda}^+(A) = \Lambda \sum_{e_i > 0} e_i + \lambda \sum_{e_i < 0} e_i,$$

where  $A$  is a matrix of order  $N$  that belongs to  $\mathcal{S}_N$ , the space of all real symmetric  $N \times N$  matrices, and  $e_i = e_i(A)$ ,  $i = 1, 2, \dots, N$ , are the eigenvalues of the matrix  $A$ . We remark that all our results here obtained are valid for the equation  $(P_\varepsilon)$ , if we replace  $\mathcal{M}_{\lambda,\Lambda}^+$  by  $\mathcal{M}_{\lambda,\Lambda}^-$ , but in order to simplify our presentation, in this work we only consider the operator  $\mathcal{M}_{\lambda,\Lambda}^+$ . Notice that Pucci defined this kind operators in the sixties, see [29]. Let us recall, from [6], the following definitions

**Definition 1.1.** *Let  $\Theta$  be a open subset of  $\mathbb{R}^N$ . We say that a continuous function  $u$  in  $\Theta$  is a viscosity subsolution (resp. viscosity supersolution) of  $(P_\varepsilon)$  in  $\Theta$ , when the following condition holds: if  $x_0 \in \Theta$ ,  $\varphi \in C^2(\Theta)$  and  $u - \varphi$  has a local maximum at  $x_0$  then*

$$-\varepsilon^2 \mathcal{M}_{\lambda,\Lambda}^+(D^2\varphi)(x_0) \leq f(x_0, u(x_0))$$

*(resp. if  $u - \varphi$  has a local minimum at  $x_0$  then  $-\varepsilon^2 \mathcal{M}_{\lambda,\Lambda}^+(D^2\varphi)(x_0) \geq f(x_0, u(x_0))$ ). We say that  $u$  is a viscosity solution of  $(P_\varepsilon)$  when it is subsolution and supersolution.*

*We say that  $-\varepsilon^2 \mathcal{M}_{\lambda,\Lambda}^+(D^2u)(x) \leq$  (resp.  $\geq, =$ )  $f(x, u(x))$  in the viscosity sense in  $\Theta$  whenever  $u$  is a subsolution (resp. supersolution, solution) of  $(P_\varepsilon)$  in  $\Theta$ .*

For the Laplacian operator, that is the case when  $\lambda = \Lambda = 1$ , problems with a nonnegative nonlinearity having a zero at a positive value were first considered in [25], and there was proved the existence of two solutions through topological degree arguments in the subcritical case. Existence and behavior of a solution below the zero of  $f$  has been studied in many works, see for instance [19] and references therein, and it can be proved that this solution converges to 1 (the positive zero of the nonlinearity) as  $\varepsilon \rightarrow 0$ . Existence of a solution whose maximum is above 1 is more delicate and usually requires some hypotheses on the growth of  $f$  at infinity. In [22], was considered the  $p$ -Laplacian operator and allowed to  $f$  depended on the  $x$  variable, but only in the subcritical case, then two positive solutions were obtained to the asymptotical problem at the origin for  $\varepsilon$  small, and was proved that both solutions converge at least pointwise to 1 as  $\varepsilon \rightarrow 0$ . This behavior suggests that also for our problem, truncation procedures like those in [7, 26, 24] could be used in order to prove the existence of two solutions when are considered critical or supercritical nonlinearities. However, the pointwise convergence is not enough to guarantee a suitable control on the  $L^\infty$ -norm of the solutions. Assuming that  $\Omega$  is convex and  $f$  is independent on the  $x$  variable, for the  $p$ -Laplacian operator in [23] was proved the existence of two positive solutions for small values of  $\varepsilon$  without imposing conditions on  $f$  at infinity.

In this work, without imposing convexity on the domain  $\Omega$  and under some local hypotheses on  $f$ , we will show the existence of at least two positive solutions for  $\varepsilon$  small, even in the case when  $f$  depends on the  $x$  variable, but without restrictions on the growth of the nonlinearity at infinity. In this way, our result generalizes the preceding ones.

In order to state our first result, we introduce the following critical exponent:

$$p^+ := \frac{\tilde{N}}{\tilde{N} - 2} \quad \text{with} \quad \tilde{N} = \frac{\lambda}{\Lambda}(N - 1) + 1.$$

In the case  $\lambda = \Lambda = 1$ , this exponent corresponds to the result of [20], where non existence of positive solution holds for

$$\Delta u + u^p \leq 0,$$

if  $p \leq p^+$ . An extension of these results to Pucci's operators was done in [11], and a generalize to more general operators, in [4, 15, 16, 2]. Here, in the proof of the Liouville type theorem (see Theorem 1.2 below), we use some ideas given in [16] related with a generalization of the arguments in [11].

Now, we are in position to state precisely our assumptions on  $f$ :

(F<sub>1</sub>)  $f : \bar{\Omega} \times [0, +\infty) \rightarrow [0, +\infty)$  is a continuous function and  $f(x, \cdot)$  is locally Lipschitz in  $(0, \infty)$  for all  $x \in \bar{\Omega}$ ,  $f(x, 0) = f(x, 1) = 0$  and  $f(x, t) > 0$  for  $t \notin \{0, 1\}$ .

(F<sub>2</sub>)  $\liminf_{t \rightarrow 0^+} \frac{f(x, t)}{t} = 1$  uniformly for  $x \in \bar{\Omega}$ .

(F<sub>3</sub>) There exist a continuous function  $a : \bar{\Omega} \rightarrow (0, \infty)$  and  $\sigma \in (1, \tilde{N}/(\tilde{N} - 2))$  such that

$$\lim_{t \rightarrow 1} \frac{f(x, t)}{|t - 1|^\sigma} = a(x).$$

(F<sub>4</sub>) There exist  $k > 0$  and  $T > 1$  such that the map  $t \mapsto f(x, t) + kt$  is increasing for  $t \in [0, T]$  and  $x \in \Omega$ .

A simple model of a function that verifies our assumptions is  $f(t) = t|1 - t|^\sigma e^t$ , while a more complex model is  $f(x, t) = (|x|t^p + \log(t + 1))e^t|1 - t|^\sigma$ , with  $p > 1$ .

Our result is the following

**Theorem 1.1.** *Assume that  $\Omega$  is a bounded smooth domain. Then, under the hypotheses (F<sub>1</sub>) through (F<sub>4</sub>), there exists  $\varepsilon^* > 0$  such that the problem  $(P_\varepsilon)$  has at least two viscosity positive solutions  $u_{1,\varepsilon}$ ,  $u_{2,\varepsilon}$ , for  $0 < \varepsilon < \varepsilon^*$ .*

*Moreover, these solutions satisfy  $\|u_{1,\varepsilon}\|_\infty \rightarrow 1^-$  and  $\|u_{2,\varepsilon}\|_\infty \rightarrow 1^+$ , as  $\varepsilon \rightarrow 0$ .*

An important tool used in the proof of the preceding results is the following Liouville type theorem for a nonnegative function with zeros.

**Theorem 1.2.** *Let  $f : [0, \infty) \rightarrow [0, \infty)$  be a continuous function satisfying the following three assumptions:*

(f<sub>1</sub>)  $f(t) = 0$  if  $t = 0$  or  $t = 1$ , and  $f(t) > 0$  if  $t \neq 1$ ,  $t > 0$ .

(f<sub>2</sub>) There exist constants  $\gamma > 0$  and  $\sigma \in (1, \tilde{N}/(\tilde{N} - 2))$  such that  $f(t) \geq \gamma(t - 1)^\sigma$ , for  $t > 1$ .

(f<sub>3</sub>) There exists a constant  $\bar{b} > 0$  such that  $\liminf_{t \rightarrow 0^+} \frac{f(t)}{t} \geq \bar{b}$ .

*Then any bounded solution of the problem*

$$(1.1) \quad \begin{cases} -\mathcal{M}_{\lambda,\Lambda}^+(D^2w) \geq f(w) & \text{in } \mathbb{R}^N, \\ w \geq 0, \end{cases}$$

*is either the constant function  $w \equiv 0$ , or else  $w \equiv 1$ .*

We strongly believe that our method can be extended to different operators for which this sort of Liouville type theorem holds, some already quoted above, and such that a Gidas-Spruck type theorem also has been obtained.

*Remark 1.1.* As our main motivation is dealing with supercritical models at infinity, two key step in our proof are: truncation arguments and a Liouville type result, which allow us to show that the solutions of the truncate problem are effectively the solutions of the original problem for  $\varepsilon$  small enough. However, note that little modifications in the proofs of our theorems above we allow us to obtain similar results in the case that  $f$  growth like  $t^p$  at infinity, with  $p \in (1, \tilde{N}/(\tilde{N} - 2))$ , but without need of using truncation arguments and some Liouville type result, see Propositions 3.3 and 3.4. A simple model in this situation is  $f(t) = t^q|1 - t|^\sigma$ ,  $0 < q \leq 1$ .

## 2. NONLINEAR LIOUVILLE TYPE THEOREM

**Proposition 2.1.** *Let  $w$  be a viscosity solution of the inequation*

$$-\mathcal{M}_{\lambda,\Lambda}^+(D^2w) \geq f(w) \quad \text{in } \mathbb{R}^N,$$

*where  $f$  is a continuous nonnegative function. Then either  $\inf_{\mathbb{R}^N} w = -\infty$ , or  $\inf_{\mathbb{R}^N} w$  is a zero of  $f$ .*

*Proof.* Let  $U(r) = \inf_{|x|=r} w(x)$ . Suppose by contradiction that  $\inf_{\mathbb{R}^N} w = M \in \mathbb{R}$  with  $f(M) > 0$ . Let  $M_0(r) = \inf_{|x| \leq r} w(x)$ . Then we must have that  $M_0(r) \rightarrow M^+$  as  $r \rightarrow +\infty$ . By the continuity of  $f$ , for a suitably large  $r_0$  and some  $\alpha > 0$ , one has  $f(w) \geq \alpha > 0$  provided  $M \leq w \leq M_0(r_0)$ .

We claim that  $U(r)$  is strictly decreasing for  $r > r_0$ . Indeed, if not, should be  $r_1$  and  $r_2$  satisfying  $r_0 < r_1 < r_2$  such that  $U(r_1) \leq U(r_2)$ , that is,  $w$  should have a minimum in  $\{x : |x| < r_2\}$ . Since in this case  $w$  satisfies

$$\begin{cases} -\mathcal{M}_{\lambda, \Lambda}^+(D^2w) \geq 0 & \text{in } B_{r_2}, \\ w \geq U(r_2) & \text{on } \partial B_{r_2}, \end{cases}$$

it results by the Strong Maximum Principle for viscosity solutions (see [6, Proposition 4.9]) that either  $w > U(r_2)$  in  $B_{r_2}$  or  $w \equiv U(r_2)$  in  $B_{r_2}$ ; the first possibility contradicts  $U(r_1) \leq U(r_2)$ ; however, by the definition of  $M_0$ , one has that if  $w \equiv U(r_2)$  in  $B_{r_2}$  then  $M_0(r_0) = M_0(r_2) = U(r_2)$ , and as a consequence  $-\mathcal{M}_{\lambda, \Lambda}^+(D^2w) \geq \alpha > 0$  in  $B_{r_2}$ , which is impossible for a constant function: this proves the claim that  $U(r)$  is strictly decreasing for  $r > r_0$ .

Now let

$$v(x) = \frac{1}{2\lambda N} |x|^2,$$

that is a radial solution of

$$\begin{cases} -\mathcal{M}_{\lambda, \Lambda}^-(D^2v) = -1 & \text{in } \mathbb{R}^N, \\ v \geq 0, \quad v(0) = 0. \end{cases}$$

Consider  $W = w + \delta v$ , with  $\delta > 0$ . Since  $\lim_{|x| \rightarrow \infty} v(x) = +\infty$ , then  $W$  has a global minimum. Since  $U(r)$  is strictly decreasing, we may choose  $\delta$  sufficiently small so that this minimum lies at some point  $x_0$ , with  $|x_0| > r_0$  and  $w(x_0) < M_0(r_0)$ . Hence,  $f(w(x_0)) \geq \alpha > 0$ . Now, choosing  $\delta < \alpha/2$ , from the definition of a viscosity supersolution, taking  $-\delta v$  as a test function we obtain

$$-\mathcal{M}_{\lambda, \Lambda}^+(D^2(-\delta v(x_0))) = \delta \mathcal{M}^-(D^2v(x_0)) = \delta \geq f(w(x_0)) \geq \alpha.$$

But this is a contradiction because  $\delta < \alpha/2$ . This completes the proof.  $\square$

*Proof of Theorem 1.2.* Let  $u$  be a bounded solution of (1.1). By Proposition 2.1 we have that the minimum of  $u$  must be a zero of  $f$ .

We initially suppose that  $\inf u = 0$  and let  $\eta : [0, +\infty) \rightarrow \mathbb{R}$  such that  $0 \leq \eta(r) \leq 1$ ,  $\eta \in C^\infty$ ,  $\eta$  nonincreasing,  $\eta(r) = 1$  if  $0 \leq r \leq 1/2$  and  $\eta(r) = 0$  if  $r \geq 1$ . It is obvious that there exists  $C > 0$  such that

$$-\mathcal{M}_{\lambda, \Lambda}^+(D^2\eta(|x|)) \leq C.$$

Define now  $\xi(x) = m(R/2)\eta(|x|/R)$ , where  $m(r) := \min_{|x| \leq r} u(x)$ . Then by scaling property of  $\mathcal{M}_{\lambda, \Lambda}^+$  we have

$$-\mathcal{M}_{\lambda, \Lambda}^+(D^2\xi(x)) \leq \frac{Cm(R/2)}{R^2}.$$

In addition,  $\xi(x) = 0 \leq u(x)$  if  $|x| > R$  and  $\xi(x) = m(R/2) \leq u(x)$  if  $|x| \leq R/2$ . Thus there exists a global minimum of  $u(x) - \xi(x)$  achieved in a point  $x_R$  with  $|x_R| < R$ . Note that  $u(x_R) - \xi(x_R) \leq 0$  and so  $u(x_R) \leq \xi(x_R) \leq m(R/2)$ .

If we define  $\varphi(x) := \xi(x) - \xi(x_R) + u(x_R)$  we obtain that  $\varphi$  is a test function for  $u$  at  $x_R$  and thus

$$f(u(x_R)) \leq -\mathcal{M}_{\lambda, \Lambda}^+(D^2\varphi(x_R)) = -\mathcal{M}_{\lambda, \Lambda}^+(D^2\xi(x_R)) \leq \frac{Cm(R/2)}{R^2}.$$

Now, since  $0 < u(x_R) \leq m(R/2)$  then by (f1) and (f3) there is  $R_0 > 0$  large enough so that

$$\frac{\bar{b}}{2}u(x_R) < f(u(x_R)) \leq \frac{Cm(R/2)}{R^2},$$

for any  $R > R_0$ , this implies that

$$\frac{\bar{b}}{2}m(R) \leq \frac{Cm(R/2)}{R^2} < \frac{cm(R)}{R^2},$$

which is impossible if  $m(R) \neq 0$  for all  $R > R_0$ . Then there is  $R > 0$  large enough so that  $m(R) = 0$ . Hence, since  $u$  is a viscosity supersolution of  $-\mathcal{M}_{\lambda,\Lambda}^+(D^2u) \geq 0$  in  $B_T(0)$  by applying the Strong Maximum Principle in [6], we have that  $u \equiv 0$  in  $B_T(0)$  for any  $T > R$ , i.e.  $u \equiv 0$  in  $\mathbb{R}^N$ .

Finally, in the case that  $\inf u = 1$  in  $\mathbb{R}^N$ , setting  $u_0 = u - 1$  we see that  $u_0$  is a viscosity solution of

$$\begin{cases} -\mathcal{M}_{\lambda,\Lambda}^+(D^2u_0) \geq f(u_0 + 1) & \text{in } \mathbb{R}^N, \\ u_0 \geq 0, \end{cases}$$

which satisfies  $m_{u_0}(R) := \min_{|x| \leq R} u_0(x) \rightarrow 0$  as  $R \rightarrow \infty$ . Arguing as in the previous case and using  $(f_2)$  we obtain that

$$\gamma u_0(x_R)^\sigma \leq f(u_0(x_R) + 1) \leq \frac{Cm_{u_0}(R)}{R^2},$$

which implies

$$m_{u_0}(R)R^{\frac{2}{\sigma-1}} \leq c,$$

i.e.

$$m_{u_0}(R)R^{\tilde{N}-2} \leq cR^{\tilde{N}-2-\frac{2}{\sigma-1}}.$$

But inequality above is a contradiction, because  $m_{u_0}(R)R^{\tilde{N}-2}$  is increasing (see [13, Corollary 3.1]).  $\square$

### 3. PROOF OF THEOREM 1.1

Note that by hypothesis  $(F_3)$  there exist  $R > 1$  and  $\gamma_1 > 0$  such that  $f(x, t) \geq \gamma_1|t - 1|^\sigma$  for  $t \in [1, R]$ . Without loss of generality we may assume that  $R \leq T$  from hypothesis  $(F_4)$ . Then we truncate  $f$  as follows

$$f_R(x, t) = \begin{cases} f(x, t^+), & t \leq R, \\ \frac{f(x, R)}{R^\sigma}t^\sigma, & t \geq R, \end{cases}$$

where  $t^+ = \max\{0, t\}$ . Also, without loss of generality, we may assume that

$$\liminf_{t \rightarrow 0^+} \frac{f_R(x, t)}{t} \geq 1 \text{ uniformly for } x \in \bar{\Omega}.$$

With this definition,  $f_R$  has a power growth at infinity with exponent less than  $\tilde{N}/(\tilde{N} - 2)$  and also satisfies the following properties:

$$f_R(x, t) \geq \gamma_2|t - 1|^\sigma \quad \text{for } t \geq 1$$

if  $\gamma_2 = \min \left\{ \gamma_1, \inf_{x \in \Omega} \frac{f(x, R)}{R^\sigma} \right\} > 0$ , and

$$(3.1) \quad \text{the map } t \mapsto f_R(x, t) + kt \text{ is increasing for } t \in [0, \infty],$$

where  $k$  is as in hypothesis  $(F_4)$ .

We consider then the auxiliary problem

$$(Q_{\varepsilon,\tau}) \quad \begin{cases} -\varepsilon^2 \mathcal{M}_{\lambda,\Lambda}^+(D^2u) = f_R(x, u) + \varepsilon^2 \tau u^+ & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\tau$  is a nonnegative parameter.

We remark that since  $f_R \geq 0$ , by the Strong Maximum Principle in [6] one has the nontrivial solutions of the problem  $(Q_{\varepsilon,\tau})$  are positive. Moreover, considering from now on  $\mu_1^+$  as the first positive eigenvalue of  $-\mathcal{M}_{\lambda,\Lambda}^+$  in  $\Omega$  (see [5, Proposition 1.1]), one gets that the problem  $(Q_{\varepsilon,\tau})$  has no positive solution for  $\tau > \mu_1^+$ .

Our first step will be to derive some a priori estimates for solutions of  $(Q_{\varepsilon,\tau})$ .

**Lemma 3.1.** *Under hypotheses  $(F_1)$  and  $(F_2)$ , we have given  $\tilde{\varepsilon} > 0$ , there exists a constant  $D_{\tilde{\varepsilon}}$  such that, if  $u$  is a viscosity solution of problem  $(Q_{\varepsilon,\tau})$  with  $0 < \varepsilon < \tilde{\varepsilon}$  and  $\tau \geq 0$  then*

$$\|u\|_{\infty} \leq D_{\tilde{\varepsilon}};$$

and therefore there is a positive constant  $C_{\varepsilon}$  such that

$$\|u\|_{C^{1,\alpha}(\bar{\Omega})} \leq C_{\varepsilon}.$$

*Proof.* Suppose, for sake of contradiction, that there exists a sequence  $\{(u_n, \varepsilon_n, \tau_n)\}_{n \in \mathbb{N}}$  with  $u_n$  being a positive viscosity solution of  $(Q_{\varepsilon_n, \tau_n})$ , such that  $S_n := \max_{\bar{\Omega}} u_n = u_n(x_n) \rightarrow \infty$  as  $n \rightarrow \infty$ , where  $\{x_n\} \subset \Omega$  is a sequence of points where the maximum is attained. We remark that since we are not supposing  $\tau > 0$  at this point, this sequence may not be bounded away from the boundary.

Let now  $\delta_n = \text{dist}(x_n, \partial\Omega)$  and define  $w_n(y) = S_n^{-1} u_n(A_n y + x_n)$ , where  $A_n$  will be fixed later. Hence  $w_n$  satisfies

$$(3.2) \quad -\mathcal{M}_{\lambda, \Lambda}^+(D^2 w_n(y)) = \frac{A_n^2}{\varepsilon_n^2 S_n} f_R(A_n y + x_n, S_n w_n(y)) + \tau_n A_n^2 w_n(y) \quad \text{in } \Omega_n = A_n^{-1}(\Omega - x_n)$$

and  $w_n(0) = \max w_n = 1$ .

We choose  $A_n^2 = \varepsilon_n^2 S_n^{1-\sigma} f(x_n, R)^{-1} R^\sigma$ . Since  $S_n \rightarrow \infty$ ,  $0 < \varepsilon_n < \tilde{\varepsilon}$  and  $\tau_n \leq \mu_1^+$  (because no positive solution of  $(Q_{\varepsilon,\tau})$  exists for  $\tau > \mu_1^+$ ), we conclude that  $A_n \rightarrow 0$  and  $\tau_n A_n^2 \rightarrow 0$ . Then, the right hand side of (3.2) becomes

$$\frac{R^\sigma f_R(A_n y + x_n, S_n w_n)}{f(x_n, R) S_n^\sigma} + o(1),$$

and by the continuity of  $f$  and the definition of  $f_R$ , it is bounded. By regularity results (see [32] for instance), we have that, up to subsequence,  $w_n \rightarrow w$  in compact sets of  $\mathbb{R}^N$  or  $\mathbb{R}_+^N$ , according to whether the limit of  $\delta_n/A_n$  is infinity or not, that is,  $\Omega_n$  tends to  $\mathbb{R}^N$  or to a half space.

Finally, taking the limit in (3.2), we have that  $w$  satisfies, in the viscosity sense, either

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2 w) + w^\sigma = 0 \quad \text{in } \mathbb{R}^N$$

or

$$\begin{cases} \mathcal{M}_{\lambda, \Lambda}^+(D^2 w) + w^\sigma = 0 & \text{in } \mathbb{R}^N, \\ w = 0 & \text{on } \partial\mathbb{R}_+^N. \end{cases}$$

But this contradicts the Liouville type theorem in [13, Theorem 4.1] in the case of  $\mathbb{R}^N$  and the corresponding in [30, Theorem 1.5] in the case of the half space, because

$$(3.3) \quad 1 < \sigma < \tilde{N}/(\tilde{N} - 2) < \frac{(\widetilde{N-1})}{(\widetilde{N-1}) - 2} = \frac{\tilde{N} - \frac{\Lambda}{\lambda}}{\tilde{N} - 2 - \frac{\Lambda}{\lambda}}.$$

This contradiction proves that  $\|u\|_{\infty} \leq C$  for any solution of problem  $(Q_{\varepsilon,\tau})$  with  $\varepsilon < \tilde{\varepsilon}$  and  $\tau \geq 0$ . Finally using the  $C^{1,\alpha}$  estimates we obtain that there is a positive constant  $C_{\varepsilon}$  such that

$$\|u\|_{C^{1,\alpha}(\bar{\Omega})} \leq C_{\varepsilon},$$

for some  $\alpha \in (0, 1)$  (see [32]). □

Now we look for a family of supersolutions of  $(Q_{\varepsilon,\tau})$ . For this purpose, we consider to the function  $\psi$  being the viscosity solution of

$$\begin{cases} -\mathcal{M}_{\lambda, \Lambda}^+(D^2 \psi) = 1 & \text{in } \Omega, \\ \psi = 0 & \text{on } \partial\Omega, \end{cases}$$

and  $A := \|\psi\|_{\infty}$ . For the existence of such  $\psi$  see, for example, [31].

**Lemma 3.2.** *Under hypothesis  $(F_3)$ , for any  $\varepsilon > 0$  there exist  $\tau_{\varepsilon}^*, \delta_{\varepsilon} > 0$  such that  $v_{\xi} = 1 + \xi + \frac{\delta_{\varepsilon}}{4A} \psi$  is a supersolution for  $(Q_{\varepsilon,\tau})$  for any  $\xi \in [-\delta_{\varepsilon}, \delta_{\varepsilon}/2]$ , and  $\tau \in [0, \tau_{\varepsilon}^*)$ . Moreover, we may choose  $\delta_{\varepsilon}$  as a nonincreasing function of  $\varepsilon^{-1}$ .*

*Proof.* Fixed  $\varepsilon > 0$ , by the hypothesis  $(F_3)$  we have that

$$\lim_{t \rightarrow 1} \frac{f_R(x, t)}{\varepsilon^2 |t - 1|} = 0,$$

and then there exists  $\delta > 0$  such that  $\varepsilon^{-2} f_R(x, t) < \frac{|t-1|}{8A} < \frac{\delta}{8A}$  for  $|t-1| \leq \delta$  and any  $x \in \Omega$ . Since this estimate still holds for lower values of  $\varepsilon$ , we deduce that  $\delta$  may be chosen as a nonincreasing function of  $\varepsilon^{-1}$ .

If  $\tau^* > 0$  is such that  $\tau u < \frac{\delta}{8A}$  for  $\tau \in [0, \tau^*)$ ,  $u \in (0, 1 + \delta]$ , then

$$\varepsilon^{-2} f_R(x, u) + \tau u < \frac{\delta}{4A} \quad \text{for } \tau \in [0, \tau^*), \quad u \in [1 - \delta, 1 + \delta] \text{ and } x \in \Omega.$$

If we define  $v_\xi = 1 + \xi + \frac{\delta}{4A} \psi$ , we have that  $v_\xi \in [1 - \delta, 1 + \delta]$  provided  $\tau \in [0, \tau^*)$ ,  $\xi \in [-\delta, \delta/2]$  and then

$$-\mathcal{M}_{\lambda, \Lambda}^+(D^2 v_\xi) = \frac{\delta}{4A} > \varepsilon^{-2} f_R(x, v_\xi) + \tau v_\xi,$$

which proves that  $v_\xi$  is a supersolution.  $\square$

Now we prove the existence of a first solution for  $(Q_{\varepsilon, \tau})$  via the sub- and supersolutions method. For this we need hypothesis  $(F_4)$ . For simplicity, for  $\mu_1^+$ , the first eigenvalue of  $-\mathcal{M}_{\lambda, \Lambda}^+$  in  $\Omega$ , we denote from now on by  $\varphi_1^+$  its first eigenfunction associated, which is positive in  $\Omega$ , see [5, Proposition 1.1] or [30]. Without loss of generality, we can consider  $\|\varphi_1^+\|_\infty = 1$ .

**Proposition 3.3.** *If hypotheses  $(F_1) - (F_4)$  hold, then problem  $(Q_{\varepsilon, \tau})$  has a positive solution  $u_{1, \varepsilon, \tau} < 1$  for  $0 < \varepsilon < (1/\mu_1^+)^{1/2}$  and  $0 \leq \tau < \tau_\varepsilon^*$ . Moreover, the following property holds: given  $0 < \bar{\varepsilon} < (1/\mu_1^+)^{1/2}$  there exists  $\rho > 0$  such that  $\rho \varphi_1^+ \leq u_{1, \varepsilon, \tau} < 1$  for any  $0 < \varepsilon < \bar{\varepsilon}$  and  $\tau \in [0, \tau_\varepsilon^*)$ .*

*Proof.* In a standard way, using  $(F_2)$ , we may find a  $\rho > 0$  (as small as desired) such that  $f_R(x, t) > \varepsilon^2 \mu_1^+ t$  for any  $t \in (0, \rho)$ , and  $0 < \varepsilon < \bar{\varepsilon} < (1/\mu_1^+)^{1/2}$ ; then  $\rho \varphi_1^+$  is a subsolution to problem  $(Q_{\varepsilon, \tau})$  for any  $\tau \geq 0$  and  $0 < \varepsilon < \bar{\varepsilon}$ .

For  $\tau \in [0, \tau_\varepsilon^*)$ , we have the supersolution  $v_{-\delta_\varepsilon} < 1$  from Lemma 3.2; since  $\delta_\varepsilon$  is not increasing in  $\varepsilon^{-1}$ , we may choose  $\rho$  such that  $\rho \varphi_1^+ < v_{-\delta_\varepsilon/2}$  for any  $0 < \varepsilon < \bar{\varepsilon}$ . Denote now by  $X$  the Banach space of  $\mathcal{C}^{1, \alpha}$ -functions on  $\bar{\Omega}$  which are 0 on  $\partial\Omega$ , endowed with the usual  $\mathcal{C}^{1, \alpha}$ -norm. Also, we will write  $u \ll v$  to say that  $u < v$  in  $\Omega$  and  $\frac{\partial u}{\partial \nu} > \frac{\partial v}{\partial \nu}$  on  $\partial\Omega$ , where  $\nu$  denotes the unitary outward normal vector to  $\partial\Omega$ . Let  $k$  be as in (3.1) and  $K_\tau : X \rightarrow X$  be defined as follows:  $K_\tau v = u$ , where  $u$  is the unique solution of the Dirichlet problem

$$\begin{cases} -\varepsilon^2 \mathcal{M}_{\lambda, \Lambda}^+(D^2 u) + ku = f_R(x, v) + (k + \varepsilon^2 \tau)v & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

the mapping  $K_\tau$  so defined is compact (see [17] for instance). Setting  $D := \{u \in X : \rho \varphi_1^+ \leq u \leq v_{-\delta_\varepsilon/2}\}$ , we have that  $K_\tau : D \rightarrow D$  is an increasing map. Thus, using the monotone iteration method (see [3] for instance), we obtain a solution  $u_{1, \varepsilon, \tau}$  with the claimed properties.  $\square$

Now, we work with  $\tau > 0$  and we show that a second solution exists: we will apply a topological degree argument, adapting a result obtained by de Figueiredo and Lions in [18], see also [23] for the  $p$ -Laplacian case.

**Proposition 3.4.** *Assume hypotheses as Proposition 3.3. If  $0 < \varepsilon < (1/\mu_1^+)^{1/2}$  and  $\tau_0 \in (0, \tau_\varepsilon^*)$ , then  $(Q_{\varepsilon, \tau_0})$  has a second positive solution  $u_{2, \varepsilon, \tau_0}$ . Moreover  $\|u_{2, \varepsilon, \tau_0}\|_\infty > 1$ .*

*Proof.* Let us fix  $0 < \varepsilon < (1/\mu_1^+)^{1/2}$  and arguing as in the proof of Proposition 3.3, we consider the bounded open set

$$\mathcal{O} = \{u \in X : \|u\|_X < C_\varepsilon + B_\varepsilon + 1, u \gg \rho \varphi_1^+\},$$

where  $C_\varepsilon, B_\varepsilon > 0$  will be chosen below (see in (3.4) and (3.6), respectively) and  $\rho > 0$  is as in the proof of Proposition 3.3, so that  $\rho \varphi_1^+ < 1$  and it is a strict subsolution for all problems  $(Q_{\varepsilon, \tau})$ ,  $\tau \geq 0$  (in particular  $\varepsilon^2 \mu_1^+ (\rho \varphi_1^+) < f_R(x, \rho \varphi_1^+)$ ).



We need that  $0 \notin (I - K_\tau)(\partial\mathcal{O})$  (i.e., no solution of  $(Q_{\varepsilon,\tau})$  lies on  $\partial\mathcal{O}$ ), so that the degree  $\deg(I - K_\tau, \mathcal{O}, 0)$  will be well defined and independent on  $\tau$ . To obtain this we consider  $C_\varepsilon$  as in Lemma 3.1, so that

$$(3.4) \quad \|u\|_X \leq C_\varepsilon$$

for all possible solutions of  $(Q_{\varepsilon,\tau})$  with  $\tau \geq 0$ .

Then, we claim that any solution  $u$  of  $(Q_{\varepsilon,\tau})$  such that  $u \geq \rho\varphi_1^+$  in  $\Omega$  satisfies  $u \gg \rho\varphi_1^+$  (and then it is not on  $\partial\mathcal{O}$ ). Actually, we have

$$(3.5) \quad \begin{cases} -\varepsilon^2 \mathcal{M}_{\lambda,\Lambda}^+(D^2u) + ku = f_R(x, u) + (k + \varepsilon^2\tau)u & \text{in } \Omega, \\ -\varepsilon^2 \mathcal{M}_{\lambda,\Lambda}^+(D^2(\rho\varphi_1^+)) + k\rho\varphi_1^+ = \varepsilon^2\mu_1^+\rho\varphi_1^+ + (k + \varepsilon^2\tau)\rho\varphi_1^+ & \text{in } \Omega, \end{cases}$$

by hypothesis  $(F_4)$ , and since  $u \geq \rho\varphi_1^+$ , we have

$$f_R(x, u) + (k + \varepsilon^2\tau)u \geq f_R(x, \rho\varphi_1^+) + (k + \varepsilon^2\tau)\rho\varphi_1^+,$$

and then a strict inequality holds between the (continuous) right hand sides of (3.5). Thus, the claim is proved.

By the above computations, we obtain that

$$\deg(I - K_\tau, \mathcal{O}, 0) = 0 \quad \text{for any } \tau > 0,$$

since  $(Q_{\varepsilon,\tau})$  has no solutions for  $\tau > \mu_1^+$ .

At this point we fix  $\tau = \tau_0$ , we consider the supersolution  $\phi := v_{\xi=0} > 1$  from Lemma 3.2, and we assume that no solution of  $(Q_{\varepsilon,\tau_0})$  touches it, otherwise such a solution would satisfy the claim and the proposition would be true. Using the  $\mathcal{C}^{1,\alpha}$  estimate in [32] we obtain that we may choose the constant  $B_\varepsilon > 0$  such that

$$(3.6) \quad \|K_\tau v\|_X \leq B_\varepsilon, \quad \forall v \in X : 0 \leq v \leq \phi;$$

we consider the open subset of  $\mathcal{O}$

$$\mathcal{O}' = \{u \in \mathcal{O} : u < \phi \text{ in } \Omega\}$$

and we claim that  $\deg(I - K_{\tau_0}, \mathcal{O}', 0) = 1$ .

Observe that  $K_{\tau_0}$  maps  $\overline{\mathcal{O}'}$  into  $\overline{\mathcal{O}'}$ . Indeed, if  $v \in \overline{\mathcal{O}'}$ , then  $\|K_{\tau_0}v\|_X \leq B_\varepsilon$  by (3.6), and if we consider  $u = K_{\tau_0}v$  we have

$$\begin{cases} -\varepsilon^2 \mathcal{M}_{\lambda,\Lambda}^+(D^2\phi) + k\phi \geq f_R(x, \phi) + (k + \varepsilon^2\tau_0)\phi, \\ -\varepsilon^2 \mathcal{M}_{\lambda,\Lambda}^+(D^2u) + u = f_R(x, v) + (k + \varepsilon^2\tau_0)v, \\ -\varepsilon^2 \mathcal{M}_{\lambda,\Lambda}^+(D^2(\rho\varphi_1^+)) + k(\rho\varphi_1^+) = \varepsilon^2\mu_1^+(\rho\varphi_1^+) + (k + \varepsilon^2\tau_0)(\rho\varphi_1^+), \end{cases}$$

then, since  $\rho\varphi_1^+ \leq v \leq \phi$ , the comparison principle implies that  $\rho\varphi_1^+ \leq K_{\tau_0}v \leq \phi$ .

Now, let  $u_0 \in \mathcal{O}'$  and consider the constant mapping  $C : \overline{\mathcal{O}'}$  defined by  $C(u) = u_0$ . So, one obtains that  $I - \mu K_{\tau_0}(v) - (1 - \mu)u_0$ ,  $\mu \in [0, 1]$ , is a homotopy between  $I - K_{\tau_0}$  and  $I - C$  in  $\overline{\mathcal{O}'}$  without zeros on  $\partial\mathcal{O}'$ : in fact, if  $v \in \partial\mathcal{O}'$  then (since  $\mathcal{O}'$  is convex)  $\mu K_{\tau_0}(v) + (1 - \mu)u_0 \in \mathcal{O}'$  for  $\mu \neq 1$ , and then it is different from  $v$ , while for  $\mu = 1$  we have  $v \neq K_{\tau_0}(v)$  since we are assuming that no solution touches  $\phi$ .

Hence  $\deg(I - K_{\tau_0}, \mathcal{O}', 0) = \deg(I - C, \mathcal{O}', 0) = 1$ , as we claimed.

Then, applying the excision property, it follows that  $\deg(I - K_{\tau_0}, \mathcal{O} \setminus \overline{\mathcal{O}'}, 0) = -1$ , so  $(Q_{\varepsilon,\tau_0})$  has a solution  $u_2 \in \mathcal{O} \setminus \overline{\mathcal{O}'}$ ; in particular,  $u_2(x_0) > \phi(x_0) > 1$  in some point  $x_0 \in \Omega$ , since otherwise it would be on  $\partial\mathcal{O}'$ , and then  $u_2$  is distinct from  $u_{1,\varepsilon,\tau_0}$  from Proposition 3.3.  $\square$

Now, we will obtain a solution for  $(Q_{\varepsilon,0})$  as the limit of the solutions obtained in the previous Proposition.

**Lemma 3.5.** *Assume hypotheses as propositions 3.3-3.4. Then, given  $0 < \varepsilon < (1/\mu_1^+)^{1/2}$ , there exists a solution  $u_{2,\varepsilon,0}$  for the problem  $(Q_{\varepsilon,0})$ , which satisfies  $\|u_{2,\varepsilon,0}\|_\infty \geq 1$ .*



*Proof.* Given  $0 < \varepsilon < (1/\mu_1^+)^{1/2}$  we will consider a sequence  $\tau_n \rightarrow 0$  and we will focus on the solution  $u_n := u_{2,\varepsilon,\tau_n}$  from Proposition 3.4, so that we know that  $\|u_n\|_\infty > 1$ .

By Lemma 3.1, we have a uniform bound for  $\|u_n\|_{\mathcal{C}^{1,\alpha}}$  for some  $\alpha \in (0, 1)$ . Then, up to a subsequence,  $u_n \rightarrow u$  in  $\mathcal{C}^1$ , where  $u$  is a nonnegative viscosity solution of  $(Q_{\varepsilon,0})$ . From  $\|u_n\|_\infty > 1$  we obtain  $\|u\|_\infty \geq 1$ . Thus  $u$  is nontrivial and then positive.  $\square$

The following Lemma is based in a result due to Quaas and Sirakov (see [30, Theorem 3.2]) which relates the existence of a nonnegative solution in  $\mathbb{R}_+^N$  with the existence of a positive solution in  $\mathbb{R}^{N-1}$ . This result will allow us to show that the maximum of the solutions are below of  $R$ , the parameter of truncation, for  $\varepsilon \neq 0$  small enough.

**Lemma 3.6.** *The solutions  $u_{2,\varepsilon,0}$  from Lemma 3.5 satisfy  $\|u_{2,\varepsilon,0}\|_\infty \rightarrow 1$  as  $\varepsilon \rightarrow 0$ . In particular, there exists  $\varepsilon^* > 0$  such that if  $0 < \varepsilon < \varepsilon^*$  then  $\|u_{2,\varepsilon,0}\|_\infty \leq R$ .*

*Proof.* Given  $\eta > 1$ , suppose by contradiction that there exists a sequence  $\varepsilon_n \rightarrow 0^+$  such that the corresponding solutions  $u_n := u_{2,\varepsilon_n,0}$  satisfy  $\|u_n\|_\infty > \eta$ , in particular there exists a sequence  $x_n \in \Omega$  such that  $d_n := \text{dist}(x_n, \partial\Omega)$  and  $u_n(x_n) = \|u_n\|_\infty > \eta$ .

Letting  $w_n(x) = u_n(x_n + \varepsilon_n x)$  we see that  $w_n$  satisfies

$$-\mathcal{M}_{\lambda,\Lambda}^+(D^2 w_n)(x) = f_R(x_n + \varepsilon_n x, w_n) \quad \text{in } B(0, d_n \varepsilon_n^{-1})$$

and  $w_n(0) = u_n(x_n)$ .

As in the proof of Lemma 3.1, we obtain (since  $w_n$  is bounded in  $L^\infty$ ) also a uniform bound in the  $\mathcal{C}^{1,\alpha}$  norm in compact sets, for some  $\alpha \in (0, 1)$ . Then, up to a subsequence,  $w_n \rightarrow w$  in the  $\mathcal{C}^1$  norm in compact sets and  $x_n \rightarrow x_0$  in  $\bar{\Omega}$ , where now  $w$  is a  $\mathcal{C}^1$  function defined in  $\mathbb{R}^N$  or in a half space. Thus,  $w$  is a  $\mathcal{C}$ -viscosity solution of the problem

$$\begin{cases} -\mathcal{M}_{\lambda,\Lambda}^+(D^2 w) = f_R(x_0, w), \\ w \geq 0. \end{cases}$$

in  $\mathbb{R}^N$  or in an half space. If such  $w$  solves the problem in  $\mathbb{R}^N$ , then according to Theorem 1.2 we conclude that either  $w \equiv 0$  or  $w \equiv 1$ . On the other hand, if such  $w$  solves the problem above in an half space, then from Theorem [30, Theorem 3.2], (3.3) and again Theorem 1.2, we conclude the same result for  $w$ .

In any case, this contradicts the fact that  $w_n(0) = u_n(x_n) > \eta > 1$ , and then the lemma is proved.  $\square$

*Proof of the Theorem 1.1.* The first solution is  $u_{1,\varepsilon,0}$ , which is the solution obtained in Proposition 3.3 for the problem  $(Q_{\varepsilon,0})$ . Indeed, note that  $u_{1,\varepsilon,0}$  verifies  $\|u_{1,\varepsilon,0}\|_\infty < 1$ . Besides, by hypothesis  $(F_1)$  and  $(F_2)$ , if  $t_\varepsilon$  is the largest real such that  $f(x, t) > \varepsilon^2 \mu_1^+ t$  for  $t \in (0, t_\varepsilon)$ , uniformly for  $x \in \bar{\Omega}$ , then  $t_\varepsilon \rightarrow 1^-$  as  $\varepsilon \rightarrow 0^+$ . Since no positive solution of  $(P_\varepsilon)$  may exist below  $t_\varepsilon$ , we deduce that  $\|u_{1,\varepsilon,0}\|_\infty \rightarrow 1^-$  as  $\varepsilon \rightarrow 0^+$ .

On the other hand, the second solution corresponds to  $u_{2,\varepsilon,0}$ , the solution to the problem  $(Q_{\varepsilon,0})$  given in Lemma 3.5, which verifies  $\|u_{2,\varepsilon,0}\|_\infty \geq 1$ . Besides, by Lemma 3.6,  $\|u_{2,\varepsilon,0}\|_\infty \leq R$  for  $\varepsilon > 0$  small, where  $R$  is the parameter of truncation, and  $\|u_{2,\varepsilon,0}\|_\infty \rightarrow 1^+$  as  $\varepsilon \rightarrow 0^+$ .  $\square$

### Acknowledgements

S.A. was partially supported by USM Grant No. 121002, L.I. was partially supported by FONDECYT No. 11080203 and Convenio de desempe\~no UTA–MINEDUC and A.Q. was partially supported by Fondecyt Grant No. 1070264 and CAPDE, Anillo ACT-125. Besides, the three authors were partially supported by Programa Basal, CMM, U. de Chile.

### REFERENCES

- [1] A. Allendes, A. Quaas, *Multiplicity results for extremal operators through bifurcation*, Discrete Contin. Dyn. Syst. **29** (2011), 51–65.
- [2] S. Armstrong, B. Sirakov, *Sharp Liouville results for Fully Nonlinear equations with power-growth nonlinearities*, preprint.

- [3] H. Amann, *Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces*, SIAM Rev. **18** (1976), 620–709.
- [4] I. Birindelli, F. Demengel, *Comparison principle and Liouville type results for singular fully nonlinear operators*, Ann. Fac. Sci. Toulouse Math. (6) **13** (2004), 261–287.
- [5] J. Busca, M. Esteban, A. Quaas, *Nonlinear Eigenvalues and Bifurcation problems for Pucci's Operator*, Ann. Inst. H. Poincaré Anal. Non Linéaire **22** (2005), 187–206.
- [6] L. Caffarelli, X. Cabré, *Fully nonlinear elliptic equations*, American Mathematical Society Colloquium Publications, 43. American Mathematical Society, Providence, RI, 1995.
- [7] S. Chen, S. Li, *On a nonlinear elliptic eigenvalue problem*, J. Math. Anal. Appl. **307** (2005), 691–698.
- [8] L.A. Caffarelli, M.G. Crandall, M. Kocan, A. Świech, *On viscosity solutions of fully nonlinear equations with measurable ingredients*, Comm. Pure Appl. Math. **49** (1996), 365–397.
- [9] F. Charro, E. Colorado, I. Peral, *Multiplicity of solutions to uniformly elliptic fully nonlinear equations with concave-convex right-hand side*, J. Differential Equations **246** (2009), 4221–4248.
- [10] J. Cheng, W. Li, *Nonnegative radial solutions to a class of elliptic equations involving the Pucci's extremal operator*, J. Math. Anal. Appl. **341** (2008), 386–397.
- [11] M. Crandall, P.-L. Lions, *Viscosity solutions of Hamilton Jacobi equations*, Trans. Amer. Math. Soc. **277** (1983), 1–42.
- [12] M. Crandall, H. Ishii, P.L. Lions, *Users guide to viscosity solutions of second order partial differential equations*, Bull. Amer. Math. Soc. (N.S.) **27** (1992), 1–67.
- [13] A. Cutri, F. Leoni, *On the Liouville property for fully nonlinear equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire **17** (2000), 219–245.
- [14] P. Felmer, A. Quaas, *Positive Radial Solutions to a 'semilinear' equation involving the Pucci's operator*, J. Differential Equations **199** (2004), 376–393.
- [15] P. Felmer, A. Quaas, *Fundamental solutions and two properties of elliptic maximal and minimal operators*, Trans. Amer. Math. Soc. **361** (2009), 5721–5736.
- [16] P. Felmer, A. Quaas, *Fundamental solutions and Liouville type theorems for nonlinear integral operators*, Adv. Math. **226** (2011), 2712–2738.
- [17] P. Felmer, A. Quaas, B. Sirakov, *Landesman-Lazer type results for second order Hamilton-Jacobi-Bellman equations*, J. Funct. Anal. **258** (2010), 4154–4182.
- [18] D. G. de Figueiredo, P.-L. Lions, *On pairs of positive solutions for a class of semilinear elliptic problems*, Indiana Univ. Math. J. **34** (1985), 591–606.
- [19] J. García-Melián, J. Sabina de Lis, *Stationary profiles of degenerate problems when a parameter is large*, Differential Integral Equations **13** (2000), 1201–1232.
- [20] B. Gidas, *Symmetry properties and isolated singularities of positive solutions of nonlinear elliptic equations*, in *Nonlinear partial differential equations in engineering and applied science*. Eds. R. Sternberg, A.Kalinowski and J. Papadakis, Proceedings of the Conference, Kingston, R.I. 1979, Lect. Notes on Pure Appl. Math., 54, Decker, New York 1980, 255–273.
- [21] B. Gidas, J. Spruck, *Global and local behavior of positive solutions of nonlinear elliptic equations*, Comm. Pure Appl. Math. **34** (1981), 525–598.
- [22] L. Iturriaga, E. Massa, J. Sanchez, P. Ubilla, *Positive solutions for the  $p$ -Laplacian with a nonlinear term with zeros*, J. Differential Equations **248** (2010), 309–327.
- [23] L. Iturriaga, S. Lorca, E. Massa, *Positive solutions for the  $p$ -Laplacian involving critical and supercritical nonlinearities with zeros*, Ann. Inst. H. Poincaré Anal. Non Linéaire **27** (2010), 763–771.
- [24] L. Iturriaga, S. Lorca, M. Montenegro, *Existence of solutions to quasilinear elliptic equations with singular weights*, Adv. Nonlinear Stud. **10** (2010), 109–120.
- [25] P.-L. Lions, *On the existence of positive solutions of semilinear elliptic equations*, SIAM Rev. **24** (1982), 441–467.
- [26] S. Lorca, P. Ubilla, *Partial differential equations involving subcritical, critical and supercritical nonlinearities*, Nonlinear Anal. **56** (2004), 119–131.
- [27] W.-M. Ni, I. Takagi, *On the shape of least energy solution to a semilinear Neumann problem*, Comm. Pure Appl. Math. **41** (1991), 819–851.
- [28] W.-M. Ni, I. Takagi, *Locating the peaks of least energy solutions to a semilinear Neumann problem*, Duke Math. J. **70** (1993), 247–281.
- [29] C. Pucci, *Operatori ellittici estremanti*, Ann. Mat. Pura Appl. (4) **72** (1966), 141–170.
- [30] A. Quaas, B. Sirakov, *Existence results for nonproper elliptic equations involving the Pucci operator*, Comm. Partial Differential Equations **31** (2006), 987–1003.
- [31] A. Świech,  *$W^{1,p}$ -interior estimates for solutions of fully nonlinear, uniformly elliptic equations*. Adv. Differential Equations **2** (1997), 1005–1027.
- [32] N. Winter,  *$W^{2,p}$  and  $W^{1,p}$ -estimates at the boundary for solutions of fully nonlinear uniformly elliptic equations*, Z. Anal. Anwend. **28** (2009), 129–164.

*E-mail address:* `salomon.alarcon@usm.cl`

*E-mail address:* `leonelo.iturriaga@usm.cl`

*E-mail address:* `alexander.quaas@usm.cl`